

Not as pathogenic as we thought?

The Queensland *Staphylococcus lugdunensis* Bacteraemia audit.

Kanthi Vemuri^{1,5}, Madan Panda^{1,4}, Robert Horvath^{1,2,4,5}, Karen Hay^{3,4}, Graeme Nimmo^{2,4,5}, and on behalf of the ieQ Collaborative.

1. Internal Medicine, The Prince Charles Hospital, Brisbane, Queensland, Australia; 2. Pathology Queensland, Brisbane, Australia; 3. QIMR Berghofer Research Institute
 4. Faculty of Medicine, The University of Queensland, Australia 5. infective endocarditis Queensland [ieQ], The Prince Charles Hospital, Queensland, Australia.

Introduction

Staphylococcus lugdunensis has a reputation of being perhaps the most pathogenic coagulase negative *Staphylococcus*, with a well documented predilection for endocarditis^{1,2,3}. It is, however, often dismissed as a contaminant.

Given the pathogenic potential and wide spectrum of clinical disease associated with *S.lugdunensis*, there is a need to study and clarify the significance of the isolates.

Aim

To determine the prevalence of infective endocarditis, true bacteraemias and contaminants amongst patients in which *S.lugdunensis* was identified in blood cultures, with reference to the number of positive isolates and clinical risk factors.

Methods

- We performed a retrospective audit of all *S.lugdunensis* bacteraemias identified in the laboratory information system of Pathology Queensland.
- 305 bacteraemias were identified, across 34 Queensland hospitals, between 1998-2017.
- An audit questionnaire was sent to all centres with 2 or more cases for clinical data to ascertain patient characteristics, risk factors for infective endocarditis and comorbidities (see *acknowledgements*).
- Based on the available information from the survey and laboratory test results, the Duke criteria were applied to establish the classification of patients into the following categories:
 - IE confirmed based on Duke criteria with *S.lugdunensis* as a causative agent
 - Possible IE based on Duke criteria with *S.lugdunensis* as a causative agent
 - Blood stream infection (BSI) with known non-cardiac source
 - BSI with unknown source
 - Contaminants


Results

- A total of 255 completed surveys have been received to date. Of these, 25 patients had definite or probable IE, giving an incidence of 10.6% amongst patients with positive *S.lugdunensis* blood cultures.
- 80% of the isolates were judged as contaminants if only one set positive, and only 2% if multiple isolates positive.
- Preliminary data of the 19 year audit suggests that the incidence of endocarditis was low (1.03% or 2/195) if only a single blood culture (BC) set was positive, but the incidence was higher if 2 (22.5% or 7/31) or more (69.2% or 18/26) were positive.
- In regards to Blood Stream Infections of a non-cardiac source, incidence was 17.7% (or 27/193) if only a single BC set was positive, and increased if 2 (58% or 18/31) or more (30.8% or 8/26) sets were positive. Data suggests that pre-existing central vascular catheters were a risk factor (30% or 16/53) for significant BSI.
- The Odds Ratio for IE if pre-existing cardiac structural abnormalities were present was 6.6 (95% CI 2.3- 9.3 p0.001).
- We also observed on review of the first 207 surveys that the introduction of MALDI-TOF* (in 2013) resulted in a 4-fold overall rise in identification of *S.lugdunensis* and a 5-fold increase in contaminants with an affiliated 2-3-fold increase in IE and significant bacteraemia.


*MALDI-TOF: Matrix Assisted Laser Desorption Ionization – Time of Flight mass spectrometry.

Enquiries: Please email Dr Kanthi Vemuri at Kanthi.Vemuri@health.qld.gov.au

Distribution of patients by diagnosis and number of positive sets



Effect on the number of positive isolates with the introduction of MALDI-TOF* in 2013 (n=207)



Conclusions

- 255 (83%) cases have so far been recorded in our audit, which we believe is the largest series of *S.lugdunensis* bacteraemia and *S.lugdunensis* IE to our knowledge.
- 27 endocarditis episodes (19 definite, 6 possible on Duke Criteria and 2 device (ICD) infections) were observed of which 92% (25/27) had 2 or more positive blood cultures.
- 53 Blood stream infection – non cardiac episodes were observed of which 49% (26/53) had 2 or more positive blood cultures.
- The significant difference in relative risk of deep seated infection with escalating numbers of positive blood cultures highlights the importance of sequential blood cultures drawn PRIOR to antibiotic commencement, especially in those with risk factors for intravascular disease.
- MALDI-TOF identification systems have resulted in a significant increase in contaminants, and a smaller but still significant increase in detection of bacteraemias, including IE.

Acknowledgements: This project would not have been possible without the support of a great number of colleagues across Queensland. We thank:

- Enzo Binotto, Robert Norton, Damen Fagg, Kayla Ward Janath de Silva
- Rudyard Yap, Basim Nona, Barry Nolan, Hsing Lee, Stephanie Tan, Andrew Cha, Zie Loo, Kathleen Poon, Eugene Ma, Nirupa Sivathasan, David Sowden
- Briana Confield, Rachel Gadge, Liam Stapleton, Paul Chapman, Thi Aung, Michael Tresillian, Andrew Redmond, Geoffrey Playford, Neil Underwood, Marjorie Sehu, Ferris Satyaputra, Andrew Henderson, Naomi Runnegar
- Clare Nourse, Michael Whitby, Michael Thomas, Mark Beale, Thomas Conrad, Rashmi Dixit, Navin Ravendran, Kylie Alcorn, Platon Choptiany, Anne Cameron and HREC TPCH

References:

- Significance of *Staphylococcus lugdunensis* bacteraemia: report of 28 cases and review of the literature. Zinkernagel A.S. et al Infection. 2008 pp 314-21
- Staphylococcus Lugdunensis*: A rare but destructive cause of Coagulase-negative *Staphylococcus* Infective Endocarditis. Sabe MA et al. Eur Heart J Acute Cardiovasc Care 2014 p 275-280
- Clinical features, outcomes and molecular characteristics of Community and Health-care Associated *Staphylococcus Lugdunensis* infections. Yeh CF et al. J Clin Micro 2016 p 2051-7